

References For Transfer functions in Micropalaeontology

Compiled by Peter Frenzel for MIO 2, Graz 2011

1. General (selection)

- Bartlein, P.J., Prentice, I.C. & Webb, T. III (1986): Climatic response surfaces from pollen data for some eastern North American taxa. – *Journal of Biogeography*, 13: 35-57.
- Birks, H.J.B. (1995): Quantitative palaeoenvironmental reconstructions. – In: Maddy, D. & Brew, J.S. (eds.), Statistical Modelling of Quaternary Science Data. Technical Guide, 5: 161-254; Quaternary Science Association, Cambridge. [overview]**
- Braak, C.J.F. ter & Juggins, S. (1993): Weighted averaging partial least squares regression (WA-PLS): An improved method for reconstructing environmental variables from species assemblages. – *Hydrobiologia*, 1: 1-18.
- Braak, C.J.F. ter, Juggins, S., Birks, H.J.B., & Van der Voet, H. (1993): Weighted averaging partial least squares regression (WA-PLS): Definition and comparison with other methods for species-environment calibration. – In: Patil, G.P. & Rao, C.R. (eds.), Multivariate environment statistics: 519-553; Elsevier, Amsterdam.
- Braak, C.J.F. ter & Looman, C.W.N. (1986): Weighted averaging, logistic regression and the Gaussian response model. – *Vegetation*, 65: 3-11.
- Guiot, J., Beaulieu, J.L. de, Cheddadi, R., David, F., Ponel, P. & Reille, M. (1993): The climate in Western Europe during the Last Glacial-Interglacial Cycle derived from pollen and insect remains. – *Palaeogeography, Palaeoclimatology, Palaeoecology*, 103, 73-93. And its application to glacial Atlantic and Pacific Oceans. – *Quaternary Science Reviews*, 24: 951-998.
- Guiot, J. & Vernal, A. de (2007): Transfer Functions: Methods for Quantitative Paleoceanography Based on Microfossils. – In: Developments in Marine Geology, 1: 523-563. [comparison of methods]**
- Imbrie, J. & Kipp, N.G. (1971): A new micropaleontological method for quantitative paleoclimatology: Application to a late Pleistocene Caribbean core. – In: Turekian, K.K. (ed.), *The Late Cenozoic Glacial Ages*: 71-181; Yale University Press, New Haven, CT. [first paper on micropalaeontological transfer functions]
- Iversen, J. (1944): *Viscum, Hedera* and *Ilex* as climatic indicators. – *Geologiska Föreningens Förhandlingar*, 66: 463-483.
- Kucera, M., Weinelt, M., Kiefer, T., Pflaumann, U., Hayes, A., Weinelt, M., Chen, M.T., Mix, A.C., Barrows, T.T., Cortijo, E., Duptat, J., Juggins, S. & Waelbroeck, C. (2005): Reconstruction of sea-surface temperatures from assemblages of planktonic foraminifera: Multi-technique approach based on geographically constrained calibration data sets
- Malmgren, B.A. & Nordlund, A. (1996): Application of artificial neural networks to chemostratigraphy. – *Paleocenaography*, 11: 505-512.
- Racca, J.M.J., Gregory-Eaves, I., Pienitz, R. & Prairie, Y.T. (2004): Tailoring palaeolimnological diatom-based transfer functions. – *Canadian Journal of Fisheries and Aquatic Sciences*, 61: 2440-2454.

Seppä, H., Birks, H.J.B., Odland, A., Poska, A. & Veski, S. (2004): A modern pollen-climate calibration set from northern Europe: Developing and testing a tool for palaeoclimatological reconstructions. – *Journal of Biogeography*, 31: 251-267.

Waelbroeck, C., Labeyrie, L., Duplessy, J.-C., Guiot, J., Labracherie, M., Leclaire, H. & Duprat, J. (1998): Improving past sea surface temperature estimates based on planktonic fossil faunas. – *Paleoceanography*, 13: 272-283.

2. Software

Hammer, Ø., Harper, D.A.T., and P. D. Ryan, 2001. PAST: Paleontological Statistics Software Package for Education and Data Analysis. – *Palaeontologia Electronica* 4(1): 9pp. [http://palaeo-electronica.org/2001_1/past/issue1_01.htm] + http://folk.uio.no/ohammer/past/Past.exe]

Juggins, S. (2007): C2 Version 1.5 User guide. Software for ecological and palaeoecological data analysis and visualisation. – 73 pp., Newcastle University. [http://www.staff.ncl.ac.uk/Stephen.juggins]

http://www.r-project.org/ + http://servpal.cerege.fr/paleocean_o_transfer/

For additional software see Birks 1995: p. 233.

3. Ostracoda

a) Set up, development, and application

Alin, S.R. & Cohen, A.S. (2003): Lake-level history of Lake Tanganyika, East Africa, for the past 2500 years based on ostracode-inferred water-depth reconstruction. – *Palaeogeography, Palaeoclimatology, Palaeoecology*, 199, 31–49. [water depth]

Cronin, T.M. & Dowsett, H.J. (1990): A quantitative method for shallow marine paleoclimatology: Application to Pliocene deposits of the western North Atlantic Ocean. – *Marine Micropaleontology*, 16 (1/2): 117-148. [bottom water temperature]

Frenzel, P. & Viehberg, F.A. (2005): Transferfunktionen für Ostrakodenassoziationen aus holozänen Ostseesedimenten. – *Berichte des Institutes für Erdwissenschaften, Karl-Franzens-Universität Graz*, 10: 14. [salinity]

Frenzel et al. (2010): Nam Co 600 year record [water depth]

Hazel, J.E. (1988): Determining late Neogene and Quaternary paleoclimates and paleotemperature regimes using ostracodes. – In: DeDeckker, P., Colin, J.-P. & Peypouquet, J.P. (eds), *Ostracoda in the Earth Sciences*, pp. 89-102; Elsevier, Amsterdam. [water temperature]

Horne, D. J. (2007): A Mutual Temperature Range method for Quaternary palaeoclimatic analysis using European nonmarine Ostracoda. – *Quaternary Science Reviews*, 26: 1398-1415. [July and January air temperature]

Horne, D.J. & Martens, K. (2011): Calibrating a new proxy for Pleistocene climate change in southern Africa. – *Joannea Geologie und Paläontologie*, 11: 89-90. [July, January, and annual air temperatures]

Horne, D.J. & Mezquita, F. (2008): Palaeoclimatic applications of large databases: developing and testing methods of palaeotemperature reconstruction using nonmarine ostracods. – *Senckenbergiana lethaea*, 88 (1): 93-112. [July and January air temperature]

Mezquita, F., Roca, J.R., Reed, J.M. & Wansard, G. (2005): Quantifying species-environment relationships in non-marine Ostracoda for ecological and palaeoecological studies:

- Examples using Iberian data. – Palaeogeography, Palaeoclimatology, Palaeoecology, 225: 93-117. [conductivity]
- Michelson, A.V. & Park, L.E. (2011): Reevaluating the assumptions of organism-based palaeoenvironmental inference models: A case study from an ostracod metacommunity on San Salvador Island, Bahamas. – Joannea Geologie und Paläontologie, 11: 141-143. [conductivity]
- Mischke, S., Almogi-Labin, A., Ortal, R., Rosenfeld, A., Schwab, M.J. & Boomer, I. (2010): Quantitative reconstruction of lake conductivity in the Quaternary of the Near East (Israel) using ostracods. – Journal of Paleolimnology, 43: 667-688. [conductivity]
- Mischke, S., Herzschuh, U., Massmann, G. & Zhang Chenjung (2007): An ostracod-conductivity transfer function for Tibetan lakes. – Journal of Paleolimnology, 38: 509-524. [conductivity]
- Mourgiart, P. & Carbonel, P. (1994): A quantitative method of palaeolake-level reconstruction using ostracod assemblages: an example from the Bolivian Altiplano. – Hydrobiologia 288, 183–193. [water depth]
- Mourgiart, P. & Roux, M. (1990): Une approche nouvelle du problème posé par les reconstructions des paléoniveaux lacustres: utilisation d'une fonction de transfert basée sur les faunes d'ostracodes. – Géodynamique, 5 (2): 151–165. [water depth]
- Peng Ping, Zhu Liping, Frenzel, P., Ju Jianting & Wrozyyna, C. (2011): Water depth related Ostracod distribution in Pumayum Co, Southern Tibetan Plateau. – Joannea Geologie und Paläontologie, 11: 156. [water depth]
- Pérez, L., Bugja, R., Massaferro, J., Steeb, P., Geldern, R. v., Frenzel, P., Brenner, M., Scharf, B. & Schwalb, A. (2010): Post-Columbian environmental history of Lago Petén Itzá, Guatemala. – Revista Mexicana de Ciencias Geológicas, 27 (3): 490-507. [water depth]
- Viehberg, F.A. (2006): Freshwater ostracod assemblages and their relationship to environmental variables in waters from northeast Germany. – Hydrobiologia, 571: 213-224. [water temperature]
- Wirrmann, D. & Mourgiart, P. (1995): Late Quaternary Spatio-temporal Limnological Variations in the Altiplano of Bolivia and Peru. – Quaternary Research, 43: 344-354. [water depth and water chemistry]
- Wrozyyna, C., Frenzel, P., Schwalb, A., Steeb, P. & Zhu Liping (2009): Water depth related assemblage composition of Recent Ostracoda in Lake Nam Co, southern Tibetan Plateau. – Revista Española de Micropaleontología, 41 (1-2): 1-20. [water depth]
- Zhu L., Xie, M., Peng, P., Wang, J., Frenzel, P., Wrozyyna, C. & Schwalb, A. (2010): Ostracod-based environmental reconstruction over the last 8,400 years of Nam Co Lake on the Tibetan plateau. – Hydrobiologia, 648: 157-174. [water depth]

b) Application only (selection)

- Bernadout, G. & Patterson, R.T. (2011): Palaeoclimate reconstructions of the last 4000 years in the Southern Northwest territories using ostracods, and implications for the long-term viability of the Tibbitt to Contwoyo Winter Road. – Joannea Geologie und Paläontologie, 11: 21-24. [July and January air temperature]
- Cronin, T.M. (1991): Pliocene shallow water oceanography of the North Atlantic Ocean based on marine ostracodes. – Quaternary Science Reviews, 10: 175-188. [bottom water temperature]

- Daniel, T. & Frenzel, P. (2011): Micropalaeontological investigations of the hominine site Bilzingsleben – new insights into a Middle Pleistocene Travertine deposit. – *Joannea Geologie und Paläontologie*, 11: 48-49.
- Mourgiart, P., Wirrmann, D., Fournier, M. & Servant, M. (1992): Reconstruction quantitative des niveaux du petit lac Titicaca au cours de l'Holocene. – *Comptes Rendus de l'Académie des Sciences*, 315 (Série II): 875–880. [water depth]
- Mourgiart, P., Corrège, T., Wirrmann, D., Argollo, J., Montenegro, M.E., Pourchet, M. & Carbonel, P. (1998): Holocene palaeohydrology of Lake Titicaca estimated from an ostracod-based transfer function. – *Palaeogeography, Palaeoclimatology, Palaeoecology*, 143: 51-72. [water depth]
- Pérez, L., Frenzel, P., Brenner, M., Escobar., J., Hoelzmann, P., Scharf, B. & Schwalb, A. (in press): Late Quaternary (24–10 ka BP) environmental history of the Neotropical lowlands inferred from ostracodes in sediments of Lago Petén Itzá, Guatemala. – *Journal of Paleolimnology*: 16 pp. [conductivity]
- Schütt, B., Berking, J., Frechen, M., Frenzel, P., Schwalb, A. & Wrozyyna, C. (2010): Late Quaternary transition from lacustrine to a fluvio-lacustrine environment in the north-western Nam Co, Tibetan Plateau, China. – *Quaternary International*, 218: 104–117. [water depth]
- Wrozyyna, C., Frenzel, P., Steeb, P., Zhu, L.P., Van Geldern, R., Mackensen, A. & Schwalb, A. (2010): Stable isotope and ostracod species assemblage evidence for lake level changes of Nam Co, southern Tibet, during the past 600 years. – *Quaternary International*, 212: 2–13. [water depth]

4. Non Ostracoda (examples)

- Abelmann, A., Brathauer, U., Gersonde, R., Sieger, R. & Zielinski, U. (1999): A radiolarian-based transfer-function for the estimation of summer sea-surface temperatures in the Southern Ocean (Atlantic sector). – *Paleoceanography*, 14: 410-421. [Radiolaria, sea surface temperature]
- Atkinson, T.C., Briffa, K.R. & Cope, G.R. (1987): Seasonal temperatures in Britain during the past 22,000 years reconstructed using beetle remains. – *Nature*, 325: 587-593. [Coleoptera, air temperature]
- De Sèze, M.A. (2005): Transfer function between surface sediment diatom assemblages and sea-surface temperature and salinity of the Labrador Sea. – *Marine Micropaleontology*, 36: 249-267. [diatoms, sea surface temperature]
- Gasse, F. & Tekaia, F. (1983): Transfer functions for estimating paleoecological conditions (pH) from East African diatoms. – *Hydrobiologia*, 103: 85-90. [diatoms, pH]
- Gersonde, R., Crosta, X., Abelmann, A. & Armand, L. (2005): Sea-surface temperature and sea ice distribution of the Southern Ocean at the EPILOG Last Glacial Maximum – a circum-Antarctic view based on siliceous microfossil records. – *Quaternary Science Reviews*, 24: 869-896. [Radiolaria, sea surface temperature]
- Guiot, J., Berger, A.L., Munaut, A.V. & Till, C. (1983): Some new mathematical procedures in dendroclimatology with examples for Switzerland and Morocco. – *Tree-ring Bulletin*, 42: 33-48. [dendroclimatology]
- Imbrie, J. & Kipp, N.G. (1971): A new micropaleontological method for quantitative paleoclimatology: Application to a late Pleistocene Caribbean core. – In: Turekian, K.K.

- (ed.), The Late Cenozoic Glacial Ages: 71-181; Yale University Press, New Haven, CT.
[planktic Foraminifera, sea surface temperature]
- Korkhola, A. & Weckström, J. (2000): A quantitative Holocene climatic record from diatoms in Northern Fennoscandia. – *Quaternary Research*, 54: 284-294. [diatoms, temperature]
- Moine, O., Rousseau, D.D., Jolly, D. & Vianey-Liaud, M. (2002): Paleoclimatic reconstruction using mutual climatic range on terrestrial mollusks. – *Quaternary Research*, 57: 162-172. [land snails, climate]
- Molfino, B., Kipp, N.G. & Morley, J.J. (1982): Comparison of foraminiferal, coccolithophorid, and radiolarian paleotemperature equations: Assemblage coherency and estimate concordancy. – *Quaternary Research*, 17: 279-313.
- Pflaumann, U., Duprat, J., Pujol, C. & Labeyrie, L. (1996): SIMMAX: A modern analogue technique to deduce Atlantic sea surface temperatures from planktonic foraminifera in deep-sea sediments. – *Paleoceanography*, 11: 15-35. [planktic Foraminifera, sea surface temperature]
- Roux, M. (1979): Estimation des paléoclimats d'après l'écologie des foraminifères. – *Cahiers de l'Analyse des Données*, 4: 61-79. [Foraminifera, temperature]
- Vernal, A. de, Turon, J.-L. & Guijt, J. (1994): Dinoflagellate cyst distribution in high latitude environments and development of transfer function for the reconstruction of sea-surface temperature, salinity and seasonality. – *Canadian Journal of Earth Sciences*, 31: 48-62. [dinocysts, sea surface temperature, salinity, and ice cover]

Types of Transfer functions (based on associations)

slightly modified from Guiot & Vernal 2007, ostracod studies added in bold

Type of approach	Family	Sub-family	Method (and selected references)	Order
Calibration	Ordination techniques	Indirect gradient analyses	Regression on principal components (Imbrie & Kipp 1979)	Linear or quadratic
			Regression on correspondence factors (Roux 1979; Cronin & Dowsett 1990? , Mourguiaut & Carbonel 1994 , Wirrmann & Mourguiaut 1994)	Linear or unimodal
		Direct gradient analyses	Partial least Square (PLS) (Braak et al. 1993)	Linear
			Canonical correlation (Guiot et al. 1983)	Linear
			Canonical correspondence analyses (Gasse & Tekaia 1983, Alin & Cohen 2003)	Unimodal
			Weighted averaging – binomial regression (WA-BR) (Wrozyńska et al. 2009, Frenzel et al. 2010, Pérez et al. 2010, Pérez et al. in press)	Linear or quadratic
			Weighted-averaging – Partial least squares (WA-PLS) (Korkhola & Weckström 2000, Frenzel & Viehberg 2005 , Mezquita et al. 2005 , Viehberg et al. 2006 , Mischke et al. 2007, Frenzel et al. 2010, Mischke et al. 2010)	Unimodal
			Logistic regression (Braak & Looman 1986)	Unimodal
			Generalized additive model (GAM) (Gersonde et al. 2005)	Non-linear
			Artificial neural network (ANN) (Malmgren & Nordlund 1996)	Non-linear
Similarity	Presence / absence	Unconstrained	Indicator species method (ISM) (Iversen 1944)	Non-linear
			Mutual climatic range method (MCR) (Hutson 1980)	Non-linear
			Mutual ostracod temperature range method (MOTR) (Horne 2007, Horne & Mezquita 2008 , Horne & Martens 2011)	Non-linear
			Modern analogue technique (MAT) (Hutson 1980)	Non-linear
	Abundances	Constrained	MAT constrained with external variable (Guiot et al. 1993)	Non-linear
			MAT constrained geographically (SIMMAX) (Pflaumann et al. 1996)	Non-linear
		Response surface	Response surface method (Bartlein et al. 1986)	Non-linear
			Revised analogue method (RAM) (Waelbroeck et al. 1998)	Non-linear